

B-Act® WSP

Targeted protection

Indications for use

Prevention of necrotic enteritis and supporting gut health in poultry.

Mode of administration

Orally, the product is dosed via the drinking water.

Dosage

1 sachet of 100 grams treats 20.000 birds per day.

Duration

Poultry should be dosed on a minimum of 7 day regime.



Huvepharma E00D • 3a Nikolay Haytov Str., 1113 Sofia, Bulgaria
P +359 2 862 5331 • F +359 2 862 5334 • sales@huvepharma.com

Huvepharma N.V. • Uitbreidingstraat 80 • 2600 Antwerp • Belgium
P +32 3 288 1849 • F +32 3 289 7845 • customerservice@huvepharma.com


B-Act[®] WSP

Targeted protection

Practical use in proportioners

3 consecutive steps:

- 1** Add 1 sachet of 100 grams directly into 10 litres of water. Do not open the sachet. The sachet will initially float, then sink to the bottom of the bucket and dissolve within 5 minutes.
- 2** Stir the solution once the sachet has fully dissolved.
- 3** B-Act[®] is now ready to use in the drinking water system.



Good drinking-line management is advised for optimal results with B-Act[®] WSP. B-Act[®] WSP can be used concurrently with antibiotics, vaccination, water acidification or chlorination.